JOURNAL OF

Environmental Health

Dedicated to the advancement of the environmental health professional

fifteen dollars

Volume 83, No. 7 March 2021

Evaluation of Barrier Sprays to Control Mosquitoes

Field Study in a Suburban Environment

Published by the National Environmental Health Association

www.neha.org


FOOD
www.insp
maryanne@inspect2go.com



Introduction

Aedes albopictus (Skuse) is an invasive mosquito species and a competent vector of several arboviruses (e.g., dengue, chikungunya, Zika). In the absence of effective vaccines, vector control is the primary means of controlling the spread of these arboviruses (Chandel et al., 2016). Larval oviposition sites for *Ae. albopictus* are diverse, ranging from natural sites (e.g., tree holes, bromeliads) to artificial containers (e.g., discarded tires, plant pot receptacles, birdbaths) (Hawley, 1988). Reducing routine sources of water-holding containers can help reduce populations of container-ovipositing mosquitoes. Furthermore, larvicides can be applied to mosquito oviposition sites to control mosquitoes before they emerge as adults. It can be dif cult for larvicides, however, to reach cryptic oviposition sources and therefore larval control has proved to be dif cult over large urban areas (Chandel et al., 2016; Fonseca et al., 2013).

Pyriproxyfen is an insecudo

FIGURE 1

Aerial View of the Study Area



Dotted outlines represent lots included in the study for A. D. Demand CS 0.03% + Archer 0.005% (every 30 days, code DA30); B. Demand CS 0.06% + Archer 0.010% (every 60 days, code DA60); C. Demand CS 0.03% (every 30 days, code D30); and D. control. White circles indicate BG-Sentinel 2 and oviposition traps. Numbers indicate de-identified codes for house addresses.

table characteristics of adult Aedes aegypti (Linnaeus). After exposure of Ae. aegypti larvae to pyriproxyfen, fecundity and fertility were measured in blood-fed adult females (Harburguer et al., 2014). In larval groups of Ae. aegypti exposed to a dose of pyriproxyfen (0.2 g/kg pyriproxyfen fumes for 8 min), adult emergence was reduced by 40% and fecundity and fertility were significantly reduced (Harburguer et al., 2014).

Lambda-cyhalothrin is a type II pyrethroid that is used in barrier sprays for adult mosquito control. Muzari et al. (2014) used leaves treated with lambda-cyhalothrin in a laboratory bioassay against Ae. aegypti and demonstrated high (>94%) knockdown after 1 hr of exposure to lambda-cyhalothrin and 100% mortality after mosquitoes were held for 24 hr in a clean container. Demand, with active ingredient lambda-cyhalothrin, 25 g/L, was applied as a barrier spray in Australia and caused a significant decrease in mosquito populations, primarily Verallina lineata (Taylor) as measured using sweep net collections between treated and control sites (Muzari et al., 2014). A study in China evaluated Demand CS, with active ingredient lambda-cyhalothrin, 20 mg/m^2, used as a barrier spray against Ae. albopictus. In this study, human landing counts were used to assess differences in abundance of mosquitoes between treatment and control properties and a reduction of 83–98% of Ae. albopictus was observed in treatment compared with control sites (Li et al., 2010).

Many mosquito control programs do not possess the personnel and/or financial resources to consistently suppress Ae. albopictus effectively through source reduction and public education campaigns in peridomestic environments (Del Rosario et al., 2014; Faraji & Unlu, 2016). In many cases, the public turns to private pest management companies for assistance with mosquito control (e.g., barrier sprays) on their properties. Therefore, it is vital that the efficacy of different barrier spray products be evaluated.

We evaluated the extent to which different application rates and frequencies of a barrier spray containing Demand CS with Archer (active ingredient: pyriproxyfen) affects life table characteristics (fecundity, fertility, adult emergence rates) of Ae. albopictus in a suburban environment. We hypothesized that pyriproxyfen would negatively affect life table characteristics in Ae. albopictus because the mosquito would contact the IGR on foliage and potentially transfer it to multiple containers within the environment, as this mosquito is known to exhibit skip-oviposition behavior. We used a combination of field and laboratory methods to evaluate impacts on mosquito abundance and life table characteristics.

Methods

Recruitment of Participants

Our study was conducted in Pitt County in Eastern North Carolina in the Cherry Oaks neighborhood historically known for abundant Ae. albopictus populations (data not shown). Homeowners were invited to participate in the study via door-to-door invitation. If homeowners were home at the time

of the investigators' inquiry, we provided a verbal description with details of the study. For homeowners who were not home, we left a handout at their front door with contact information for investigators. Investigators conducted two to three follow-up visits until the homeowner was contacted and invited to participate in the study.

In total, 12 residences (grouped by three nearby residences for each treatment type) were targeted for recruitment in our study. Control properties were recruited and were at least 100 m from treatment properties. Participants were instructed not to carry out any insecticide treatments in their yards for the duration of the study. Barrier sprays were provided to homeowners free of charge for the duration of the study to encourage participation. The institutional review board (IRB) at East Carolina University was consulted and determined that the study did not meet the federal de nitions of research involving human participants, hence full IRB review was not required.

TABLE 1

Total Adult Mosquitoes Collected From BG-Sentinel 2 Traps

Mosquito Species	Treatment			
	Control	Demand CS 0.03% (Every 30 Days)	Demand CS 0.03% + Archer 0.005% (Every 30 Days)	Demand CS 0.06% + Archer 0.010% (Every 60 Days)
Aedes albopictus	371	193	396	392
Ae. atlanticus	311	199	22	21
Ae. canadensis	3	4	0	0
Ae. japonicus	1	1	2	1
Ae. tormentor	60	29	2	11
Ae. triseriatus	3	7	1	0
Ae. vexans	26	9	10	36
Anopheles crucians complex	4	4	3	1
An. punctipennis	118	88	70	112
An. quadrimaculatus	5	4	7	7
Culex erraticus	44	51	17	26
Cx. pipiens/ quinquefasciatus	23	16	8	9
Cx. restuans	1	0	0	0
Cx. salinarius	0	0	0	1
Orthopodomyia signifera	3	1	0	0
Psorophora ciliata	5	1	6	2
Ps. columbiae	25	29	76	60
Ps. ferox	51	106	16	103
Ps. howardii	0	0	0	1
Toxorhynchites rutilus	1	0	0	0

Treatments

Certi ed pest control operators from Clegg's Pest Control (private company with a franchise location in Greenville, North Carolina) carried out barrier sprays for the project. Properties were treated via barrier sprays (Stihl SR 200 backpack blower mister) by a Clegg's Pest Control operator as follows (Figure 1):

1) Demand CS 0.06% + Archer 0.010% (every 60 days; treatment dates of June 13, August 15, and October 17) (code DA60).
2) Demand CS 0.03% + Archer 0.005% (every 30 days; treatment dates of June 13, July 13, August 15, September 15, and October 17) (code DA30).
3) Demand CS 0.03% (every 30 days; treatment dates of June 13, July 13, August 15, September 15, and October 17) (code D30).
4) Control (not treated).

The label recommends Demand CS be applied at the 0.06% rate for residual control of mosquitoes. In our study, we used this rate at an interval of 60 days and a lower rate (0.03%) at a more frequent interval of 30 days to evaluate ef cacy. Similarly, the label recommends Archer be applied at the 0.010% rate for residual control of mosquitoes and in our study, we used this rate at an interval of 60 days and a lower rate (0.005%) at a more frequent interval of 30 days to evaluate ef cacy. Operators, following label instructions, applied 2–5 gallons of the nished solution per 305 m^2 in circular patterns to vegetation until runoff. Treatments were not conducted in high winds or misty/rainy conditions. We coordinated with the Pitt County Vector Control manager and the City of Greenville Public Works mosquito control operators to let them know of the ongoing study and requested that no insecticides be sprayed in the study area for the duration of the project.

Host-Seeking Mosquitoes

Host-seeking mosquitoes were sampled weekly May 16–October 31, 2017, using BG-Sentinel 2 traps (BioQuip) baited with human scent lure, octanol, and carbon dioxide. Carbon dioxide tanks containing regulators set to release gas at 200 ml/min (BioQuip) were af xed upright to a shepherd's hook pole and a clear vinyl tube (1/4-in. outside diameter; 1/8-in. inside diameter) was clipped to the opening of the BG-Sentinel 2 trap. Each week, a BG- Sentinel 2 trap was set at each study residence (N = 12 traps) for a 24-hr period starting at approximately 9 a.m.

BG-Sentinel 2 traps were placed in a shaded area in the approximate center of properties within the barrier zone. When traps were retrieved the next morning at approximately 9 a.m., adult mosquitoes were transported to the laboratory on ice, identi ed to species using a dichotomous key (Harrison et al., 2016), and counted using a dissecting microscope. Data for each trap were tracked in Excel according to property address and date.

Weather Monitoring

Weekly averages for temperature and precipitation were retrieved and tabulated from Weather Underground, an online source of historical weather data. Time lags of 1, 2, 3, and 4 weeks were computed and used in statistical analyses of weather variables in relation to mosquito abundance.

Data Analyses

Statistical analyses were carried out using SAS and comparisons with $p < .05$ were considered se5 wp2aysw

Assessment of Oviposition Intensity

Egg-laying intensity of container-ovipositing mosquitoes was monitored to determine whether treatments affected this measure of fecundity. We expected eggs laid in ovitraps to originate from gravid mosquitoes residing in the same yards, as well as from mosquitoes immigrating into the yards from other yards in the vicinity. Eggs were collected weekly at all 12 sites (treatment and control) by using a standard oviposition trap (i.e., black plastic 500-ml cup half lled with tap water containing an oviposition substrate of seed germination paper (2.5 x 7 cm) placed inside around the circumference and drainage holes drilled 4 cm from the lip. The cup was zip tied to the same shepherd's hook pole to which the carbon dioxide tank (for the BG-Sentinel 2 trap) was af xed.

At each property, one ovitrap was placed continuously on the ground. The oviposition substrate was replaced weekly (when BG-Sentinel 2 traps were retrieved) for the duration of the study. Each week, the oviposition substrate was transported back to the laboratory and eggs were identi ed to mosquito species (Bova et al., 2016), counted, and added to data sheets coded for each property. The water in the cups was poured into Whirl-Pak bags each week, because larvae sometimes hatched on the strips; we tracked this information, also. After emptying each cup, fresh tap water was poured into the ovitraps.

Assessment of Life Table Characteristics

After eggs on oviposition substrates were counted, egg strips were dried (to stimulate hatching) and then submerged in 450–750 ml of tap water in emergence cages (Bio-Quip). For the purposes of this study, fecundity equals the number of eggs laid per trap, although we did not track how many mosquitoes laid eggs in each trap. We kept emergence cages in an incubator at 28 °C and fed larvae liver powder ad libitum. Approximately 5 days post-hatching, the number of larvae were counted to calculate fertility (fertility equals the number of larvae per trap divided by number of eggs on ovistrips per trap) and allowed to reach adulthood. Larvae that had hatched prior to the egg strip being retrieved (i.e., collected in Whirl-Pak bags) were also counted in the life table characteristics measures. Adults (females and males) from each brood, separated by date and address, were identi ed to species, counted, and sacri ced.

Weekly Means (± SE) of *Aedes albopictus* Adults Collected in BG-Sentinel 2 Traps



Note. Red arrows indicate treatment weeks.

Week

showed a signi cantly higher mean number of Ae. albopictus per BG-Sentinel 2 trap than the other weeks of the study (df = 22; F = 2.65; p = .002); however, no differences between treatments were observed (df = 3; F = 1.06; p = .417; Figure 3).

After week 24 (when treatments had commenced), no signi cant differences were observed in the abundance of host-seeking female Ae. albopictus between treatments (df = 3; F = 0.99; p = .444). We did observe, however, differences in abundance of Ae. albopictus adults between weeks when all treatments were considered in the analyses (df = 17; F = 2.41; p = .003). When analyses were performed for each treatment type individually, analyses after week 24 indicated no signi cant differences in mean numbers of host-seeking Ae. albopictus between weeks in traps placed at DA60 lots (df = 17; F = 1.36; p = .243) or DA30 lots (df = 17; F = 1.31; p = .259). Signi cant differences, however, were observed in mean numbers of host-seeking Ae. albopictus per trap between weeks in D30 lots with the highest in week 29 (July 19 and 20) (df = 16; F = 2.58; p = .022) and also in control lots, with the highest in week 27 (July 6 and 7) and week 37 (September 14 and 15) (df = 22; F = 2.45; p = .009). Figure 3 presents this information.

The numbers of host-seeking Ae. albopictus collected in control lots (df = 17; F = 2.66; p = .002), D30 lots (df = 16; F = 2.78; p = .015), and DA60 lots (df = 17; F = 2.86; p = .012) could be predicted by average temperatures during the week of collection, but not for DA30 lots (df = 17; F = 1.31; p = .259). Likewise, rainfall during the week of collection was a positive predictor of host-seeking Ae. albopictus collected in traps in the D30 lots (df = 17; F = 3.92; p = .05).

Aedes albopictus Eggs

We collected a total of 4,423 Ae. albopictus eggs in ovitraps during the study from May 16, 2017, to November 2, 2017. The mean numbers of Ae. albopictus eggs per trap for each treatment are shown in Figure 4. Signi cant differences were observed in the abundance of Ae. albopictus eggs between treatments (df = 3; F = 4.62; p = .037), with the control lots having higher abundance compared with treatment lots. Conversely, we did not observe statistically signi cant differences in abundance of Ae. albopictus eggs between weeks (df = 19; F = 1.05; p = .412; Figure 5).

Data for ovistrips collected from the eld and mosquitoes reared in the laboratory are shown in Figures 5 and 6. Signi cant differences were observed in the mean numbers of larvae hatched per ovitrap (fertility) between treatment groups (df = 3; F = 4.32; p = .043). Significantly more larvae of all species hatched from eggs on strips collected from control lots compared with other groups. We observed a similar pattern in the mean numbers of Ae. albopictus adults (females and males that were reared in the laboratory from ovistrips collected in the eld; df = 3; F = 2.82; p = .041) and total adults of all species (df = 3; F = 4.04; p = .050) among treatment groups. Furthermore, signi cantly more adult Ae. albopictus—and adults of all species—emerged in the control group compared with other groups.

The number of

Weekly Means (± SE) of *Aedes albopictus* Eggs Collected in Ovitraps



Note. Red arrows indicate treatment dates.

2016), especially when single properties are treated but aresurrounded by untreated properties. Agreater distribution of BG-Sentinel 2 traps within the neighborhood could improve the estimation of adult mosquitoes (VanDusen et al., 2016); however, the skip oviposition behavior of Ae. albopictus increases the likelihood that these mosquitoes will oviposit in containers from neighboring yards within its ight range of <200 m. It is likely that application technique; frequency of application; scale (i.e., number of adjacent properties treated); number, size, and organic content of water-holding containers present; and other unknown factors impact ef cacy (Unlu et al, 2018). This type of barrier spray treatment using amixture of adulticide and IGR shows promise and should be evaluated further.

Truck-mounted ultralow volume application of pyriproxyfen has also been shown effective for control ofAe. albopictus (Doud et al., 2014; Unlu et al., 2017) and should be investigated further as a control method for Ae. albopictus and other container-ovipositing mosquitoes. Wenoted that foliage and brush was cut away at one of the properties within the DA30 lot during weeks 38–39 (late September, approximately 2 weeks after a treatment), which might have affected trap counts during this period. FurAe. albopictAr268 -1.2 Txtursamtemberies

couaiphich mighs Tcn

co Tc 0.14 eggontw -.yripr8 (o074 Idictus)T period. ected (tT* sible, 7 Tw bopictAfother)ictAr26o which might o 9.8ng cctive

properties, and other unknown factors that we did not assess.

Interestingly, the lowest number of hatched larvae and Ae. albopictus adults that emerged came from the D30 group, which could illustrate some degree of natural variation in Ae. albopictus abundance between lots and/or that the adulticides affected egg laying and/or hatch rates. The reason for assessing life table characteristics (i.e., fecundity, fertility) for eggs laid in the eld in the different control and treatment properties was to determine if the IGR and/or adulticide affected egg laying or hatching. While it would be dif cult to ascertain the degree to which mosquitoes from adjacent untreated properties laid eggs in our ovitraps, we see this study as a starting point for evaluating this speci c IGR-adulticide mixture used in barrier spray applications. Laboratory studies are ongoing and we will further analyze the relationship between . Wh30Vposre un bdultimosquitoes -

Li, C.X., Wang, Z.M., Dong, Y.D., Yan, T., Zhang, Y.M., Guo, X.X., Wu, M.Y., Zhao, T.Y., & Xue, R.D. (2010). Evaluation of lambda-cyhalothrin barrier spray on vegetation for control of Aedes albopictus in China. Journal of the American Mosquito Control Association, 26(3), 346–348. https://doi.org/10.2987/10-6007.1

Muzari, O.M., Adamczyk, R., Davis, J., Ritchie, S., & Devine, G. (2014). Residual effectiveness of lambda-cyhalothrin harbourage sprays against foliage-resting mosquitoes in north Queensland. Journal of Medical Entomology, 51(2), 444–449. https://doi.org/10.1603/me13141

Rhyne, M.N., & Richards, S.L. (2020). Impact of the insect growth regulator pyriproxyfen on immature development, fecundity, and fertility of Aedes albopictus. Journal of the American Mosquito Control Association, 36(1), 11–15. https://doi.org/10.2987/19-6893.1

Unlu, I., Suman, D.S., Wang, Y., Klinger, K., Faraji, A., & Gaugler, R. (2017). Effectiveness of autodissemination stations containing pyriproxyfen in reducing immature Aedes albopictus populations. Parasites & Vectors, 10(1), 139. https://doi.org/10.1186/s13071-017-2034-7

Unlu, I., Williams, G.M., Rochlin, I., Suman, D., Wang, Y.,

Introduction

Rodents (over 1,400 species worldwide) can cause damage to crops (both in the eld and in storage), forestry, nursery and ornamental plants, rangeland, cable, and irrigation pipes; they also can bite and transmit diseases to humans and other mammals. Rodents can even lead to the extinction of native ora and fauna when introduced to islands (Witmer & Eisemann, 2007). In the U.S., native species that cause signi cant damage in various regions include beavers (Castor canadensis), deer mice (Peromyscus spp.), ground squirrels (Spermophilus spp.), marmots (Marmota spp.), mountain beavers (Aplodontia rufa), pocket gophers (Thomomys spp., Geomys spp.), porcupines (Erethizon dorsatum

infected rodent. Signs and symptoms of allergic reactions from rodents can include angioedema, asthma, bronchospasm, conjunctivitis, rhinitis, urticaria—and rarely, anaphylaxis (Hesford et al., 1995; Matsui, 2013; Rankin et al., 2007; Sheehan et al., 2010; Trummer et al., 2004). In addition, the mental health impacts of rat exposure on urban residents is underappreciated (Lam et al., 2018).

The prevalence of asthma and rat infestation is high in inner cities. In one study in New York City, 13% of apartments were found to have mice present and 37% of the apartments had one or more residents with asthma (Chew et al., 2006). In a study of animal bites in Jefferson County, Alabama, 11% were due to rodents, both wild and pets (Maetz, 1979). A 2003–2006 New York City emergency department (ED) survey (Bregman & Slavinski, 2012) cataloged over 24,000 animal bites, and of these, 8.5% were from rodents (1,614 were from mice and rats, 273 from squirrels, and 173 from hamsters).

Bites and allergic reactions from rodents are also well-recognized occupational hazards in laboratory animal workers, veterinarians, and animal control of cers (Anderson et al., 1983; Kampitak & Betschel, 2016; Mann et al., 1984). In a national study of non-canine bite-related injuries in EDs in the U.S. from 2001–2010, there was an annual average of 13,707 rodent bites reported, accounting for 1.4% of reported non-canine animal-related injuries (Langley et al., 2014). Of these bites,

hospitals were combined with patients hospitalized, as both case dispositions indicate a need for a higher level of care.

We coded the narrative descriptions for the source of the bite. The source was extracted from the narrative using text string queries and then each comment line was read to verify that the correct one was coded. We replicated this veri cation twice. A sample weight was calculated for each injured per son treated at a NEISS-AIP hospital based on the inverse probability of selection of that hospital. In addition, sample weights were adjusted for nonresponse and poststrati ed to adjust for changes in the annual number of ED visits over time. Unfortunately, classi- cation of the rodent as a pet or wild animal was not available. Rates were calculated using bridged race population estimates from the U.S. Census Bureau. Injury estimates were identi ed as unstable if the national estimate was <1,200, the number of sample cases used was <20, or coef cient of variation (CV) was >30%, where CV = (SE/national estimate) x 100 (Vyrostek et al., 2002). Analyses were conducted with SAS version 9.4.

Results

From 2001–2015, an estimated 190,900 rodent bites were reported in EDs across the U.S., amounting to over 12,700 treated bites annually. Injury estimates are presented by source, demographics, month, case disposition, and affected body part. The most frequently reported rodent bites were from rats (33.6%), mice (24.1%), and squirrels (22.4%). Bites from rodents traditionally considered pets (hamsters 8.4%, guinea pigs 1.9%, and gerbils <1%) were a much smaller proportion of reported bites (Table 1). e thnt onequparber of(the patient)219.9 ( )]TJ 0 Tc 02238 Tw -1 -1.2 Td [((6.71%) wer)19.8 (e< 10yeaers of ber((7.71%) of the cyse(. lf.)]TJ -0.085 Tw T* [inerllynalf cyses(972.4%,e patiente(wer)19.8 (e tr)19.8 (eated)]TJ 0.038 Tw T* [andur)19.7 (

risk (Childs et al., 1998). Interventions that involved improved garbage management and street sanitation within a designated area reduced the likelihood of nding rats.

Natural disasters (e.g., hurricanes, earthquakes) can change the ecology of a particular area, making it more favorable for rats and other pests at a time when community services are already taxed (National Science Foundation, 2015). Rain, wind, and ooding can damage natural nesting areas of rodents, forcing them to seek higher ground and food sources inside homes, barns, and poultry and swine houses—thus potentially encountering human occupants.

Control of rodents, primarily rats and mice, is important to prevent damage of buildings and to protect the health and safety of occupants (CDC, 2006). Damage to a structure can occur when rats and mice gnaw on structural components, such as wiring, wood, and plastics. The gnawing on wire insulation can result in electrical shorts and res (Desoky, 2018a). A variety of methods can be used to manage rodent populations directly or to reduce the damage caused by rodents. Like humans, rodents require food, water, and shelter. Long-term damage mitigation and population control results generally are best achieved if a variety of methods are used (CDC, 2006, 2010; CDC & U.S. Department of Housing and Urban Development, 2006; Desoky 2018a, 2018b; Witmer, 2019).

Integrated pest management (IPM) is a comprehensive interventional approach that emphasizes the health and safety of the environment, humans, and nontarget animal species (e.g., pets, birds, and agricultural animals). Successful IPM programs require an understanding of the behavior, ecology, and activity of the target rodent pest in the environment, as well as the changes that periodically occur in the pest's environment (i.e., human-driven and climatic events) (CDC, 2006). Conducting rodent surveillance of exterior areas of residential and abandoned buildings; commercial (e.g., restaurants) and public buildings (e.g., schools); vacant lots and other public areas (e.g., parks); and agricultural production facilities is necessary to assess the conditions of infestation in a community.

A multisectoral approach includes managing rodents by preventing and treating rodent ins the 8 -insex9 ( )gity.r 8 (N(ood,)]TJ -icto [(stapublic (anuldppr)19.8ndof)sweepic ar)19.8 (

American Veterinary Medical Association. (2018). Pet ownership and demographics sourcebook. https://www.avma.org/sites/default/ les/resources/AVMA-Pet-Demographics-Executive-Summary.pdf

Anderson, L.C., Leary, S.L., & Manning, P.J. (1983). Rat-bite fever in animal research laboratory personnel. Laboratory Animal Science, 33(3), 292–294.

Bonnefoy, X., Kampen, H., & Sweeney, K. (2008). Public health signi cance of urban pests. World Health Organization.

Bregman, B., & Slavinski, S. (2012). Using emergency department data to conduct dog and animal bite surveillance in New York City, 2003–2006. Public Health Reports, 127.

CDP In

For more information or to schedule a demo,
visit www.cdpehs.com or
call CDP at (800) 888-6035.

government agency staff member, two local elected officials, and a number of community and health activists.

It was important for the researchers to understand the community issues from the different points of view of the community members interviewed. During this process, we discovered that the important organizations involved had vested interests. For example, both ETU and CFMEU representatives were supportive of their workers, with the ETU representative stating that pressure needed to be placed on AGL to assist workers without jobs to diversify through assistance with start-up companies. This statement was followed by the CFMEU representative suggesting that the Australian Labor Party had a worker retrenchments/redundancies plan for dealing with Liddell while the national CFMEU representative was sure a worker transfer plan, such as the one used in Latrobe Valley, Victoria, would be effective for Liddell workers.

On the other hand, AGL's communications representative shared with the researchers the company's renewables plan without any reference to worker needs. Taking another position, the mayor of Muswellbrook was keen to discuss what had been done to develop partnership opportunities with universities and other organizations for initiatives in UHV. The mayor commented, "When council consults with community, there is a repeated and consistent concern threaded through our feedback—where will our kids work in 20 years from now?"

Survey

In order to complement the qualitative approach, the author developed a survey during February 2018 that included closed- and open-ended questions about:

- individual and family health and well-being,
- types of assistance that might be required by those affected by the Liddell closure,
- community impacts of the Liddell closure,
- whether the Liddell closure was the beginning of a transition away from reliance on coal,
- whether a transition away from coal in UHV would have a significant impact on the life of their communities,
- types of assistance that might be required if the transition away from coal was occurring in UHV,
- whether people were interested in working with renewables, and
- demographic questions including participant age and the type of work done by the individuals and their spouses/partners.

In the survey, 10 of the questions involved a 5-point Likert scale (strongly disagree, disagree, neither agree nor disagree, agree, and strongly agree), 5 open-ended questions, and 9 closed-ended questions. After a survey draft was distributed to colleagues for review and comment, distribution of the final survey in paper form commenced in March 2018 and in electronic form in April 2018. The survey closed in late May 2018.

To obtain a wider cross section of the community, surveys were distributed at the two advertised community meetings and via street stalls in the main street, and at a local Muswellbrook pharmacy. The electronic version of the survey was completed by people working for Liddell after AGL approved its distribution and announced it to staff(distribgraphic)9iddell after AGL 0 Tw T* ter aftf the

- People worry for their children's job future, especially because young apprentices thought they would be working at Liddell. Discussion about redeployment is disappointing because there are not many diverse industries around Muswellbrook.
- Disagreement occurred about Muswellbrook's coal being the last to shut down. One viewpoint was that mining would be shut down through global sanctions because more people will go off the grid, banks will pull out of funding for mining, and the world will become "realistic."
- Government policy and planning is lacking. The community needs inspirational bipartisan leadership.
- The local community does not care about these issues. It could be ignorance, lack of education, or not wanting to know. One respondent who was a local elected official offered that she got hate mail about her support of renewables on social media, which could reflect some attitudes in UHV communities.

Focus Group Meetings

Once all the surveys had been collected, the quantitative data were analyzed statistically and the qualitative data from surveys and community meetings were examined for themes. The themes were separately identified by the chief researcher and research assistant and agreed on, resulting in a set of preliminary findings that could be discussed with the two focus groups. Notes were taken at the focus groups. For the purpose of validation, four people attended the focus group meetings held in Muswellbrook on June 14 and 20, 2018, to discuss, review, and agree with the preliminary findings, as well as add their understanding. The survey asked respondents whether they wanted to participate in the focus groups. People indicating interest were contacted by phone. In addition to scheduled participants, on June 14, a friend of a focus group participant and a Muswellbrook elected official attended; on June 20, a worker from Liddell also attended to provide his input.

Results

Demographics

A total of 98 surveys were returned (60 paper and 38 online). Of the respondents, 54% were women and 46% were men. The age of respondents ranged from <20 to >70 years (Figure 1). In regard to employment, 82% of respondents stated they were working, 11% were retired, and 5% were not working. A small number (2%) said they were concerned about losing their job or someone close to them losing their job.

Respondents shared a wide range of areas of employment, as did their spouses/partners, with the two biggest combined groups being power plant and nursing/healthcare workers (Figure 2). The majority of respondents (58%) stated they lived in Muswellbrook, with the remainder split evenly across nearby locations.

Community Uncertainty

Our researchresear th17.42d [(focua)- >7(re-0.01

would power the town. The remaining respondents' answers suggested they were ambivalent about the closure. Nearly three quarters of respondents (71%) indicated that the transition away from coal will signi cantly impact the economic and social life of UHV communities.

Health and Well-Being

A majority of respondents (55%) saw themselves and their families as being affected by the announced closure of Liddell, with most respondents being in the 51–60 age group.

tion also expressed doubts about the reliability of renewable energy to generate sufficient electricity at a good price, which also appeared linked to respondents' concerns over transferability of their skills (Figure 6).

The difficulty getting accurate information about coal and its health hazards, as well as about renewables and their capabilities, was noted by the focus groups. Additionally, because coal is an important economic factor in their community, some people had formed a denial response.

Discussion

Due to concern shown in the study findings regarding the lack of community knowledge about their health and renewable energy, the author's focus is on these aspects in the discussion, recommendations, and conclusion.

Health

It has been previously shown that toxic chemicals produced by power stations contribute to community illness, along with huge estimated health costs from coal mining and power generation per annum in UHV (Climate and Health Alliance, 2015). AGL's self-reported pollution data state that for 2016–2017, the Liddell Power Station emitted 28 kinds of pollutants, the most deadly being sulphur dioxide (33,490 tonnes), nitrogen oxides (18,627 tonnes), $PM_{2.5}$ (18.3 tonnes), in addition to emissions of carbon dioxide (8,855,569 tonnes) (Australian Government, Clean Energy Regulator, 2018; Australian Government, Department of the Environment and Energy, 2018).

Fiona Plesman, former acting general manager of Muswellbrook Shire Council, acknowledged that the levels of nitrogen oxides would be a real community concern, especially for regulators and the council, as they were 3 times the rate of global emissions (Millington, 2018). Focus group members were surprised that so many respondents believed their health would not improve (26%) or were undecided about any health improvement (46%) with the power station closure. Survey comments included, for example, "As far as I am aware only steam is emitted from the Liddell Power Station" and "There are so many mines around us that one more source of pollution won't make any difference."

A Renewable Energy Hub for Upper Hunter Valley

Power workers had the highest interest in working with the renewable energy and technology sectors because they could transfer their existing skills to alternative employment such as renewable energy, working with new start-up industries as well as the decommissioning of power stations and mines.

As stated previously, AGL is developing renewable energy sources, including some on the existing Liddell site (AGL, 2017; CFMEU, 2017). In June 2008, the Centre of Full Employment and Equity at The University of Newcastle published their policy report, A Just Transition to a Renewable Energy Economy in the Hunter Region, Australia (Bill et al., 2008). This report models and analyzes two different renewable energy scenarios for the Hunter and Wyong regions. Furthermore, it highlights that if there is a shift from coal-red power generation to a renewable energy economy, the creation of thousands of well-paid

paper, Sharing the Challenges and Opportunities of a Clean Energy Economy. The policy paper argues for navigating and managing Australia's transition to a clean energy economy (Australian Council of Trade Unions, 2016; Slezak, 2016). Global discussion on this topic is ongoing (Sweeney & Treat, 2018).

In planning for the future, collaboration and cooperation should include state and local government, representatives of chambers of commerce, relevant industry bodies, union representatives, and community groups, in line with recommendations of the Liddell Innovation Project (AGL, 2018). Respondents are supporting a broad-based, inclusive community planning and delivery approach with key players ensuring new start-up industry opportunities and retraining through TAFE, which might t well into the Liddell Innovation Project.



Communication and Education

The focus group participants identi ed the dif culty in obtaining clear, accurate, and consistent information about the health impacts of coal and the features of the various kinds of renewable energy technologies. This nding spotlighted that communication is a major issue. A communication change management program led by parties involved in the focus on transition would need to have both education and communication roles.

Limitations

This study was conducted at a time when the coal- red Liddell Power Station was under re from political sabotage. This timing could have had an in uence on the research ndings. The federal government has put pressure on Liddell to stay open using tactics such as nes or enacting laws that could force Liddell to remain open (Cox, 2019). Liddell still generates 10% of NSW's power supply despite its age, condition, and high maintenance costs. AGL is keen to replace Liddell with a combination of renewables, gas, and storage, which they argue would be cheaper than keeping Liddell open (Macdonald-Smith & Potter, 2018).

Another limitation could be the small sample size of this survey. A mixed research method using the triangulation approach to validate survey data provided evidence to support this research project (Noble & Heale, 2019). In addition, this study included a reasonable cross section of the local community (e.g., nursing/healthcare professionals, mining, retail, agriculture, viticulture, education, administration, and Liddell power workers).

Recommendations

Although mine dust is a problem that needs to be tackled, the rst concern prior to the closure of Liddell should be that it is unacceptable to allow this power station to emit nitrogen oxides at a rate much higher than international standards (Millington, 2018). Coal- red air pollution continues at the same rate despite a recent license review in January 2019 by the NSW Environment Protection Authority and population research evidence of resultant serious health problems (Ewald, 2018; Smith et al., 2013).

The second concern is a lack of awareness and understanding of the serious health effects of power station emissions. This knowledge gap was illustrated by respondents believing their health would not improve (26%) or those who were undecided about any health improvements (46%) resulting from the Liddell closure. Also, respondents were more worried about the dust produced by coal mining than the toxic emissions from Liddell. This gap needs to be addressed with an educational program focused on how mine and power station emissions affect health. The Muswellbrook Shire Council, working with NSW Department of Education, NSW Ministry of Health, and AGL, should fund and develop these health promotion and community education programs.

Conclusion

Through a mixed-methods research approach consisting of community and focus group meetings, key community personnel interviews, and a survey, the author was able to provide evidence that respondents were uncertain about the effects on their own health and that of their family. Many respondents were unclear whether the closure of Liddell would affect them; however, 71% of respondents clearly agreed that the transition away from coal would have a signi cant impact on the economic and social life of UHV communities.

There was only a small group of respondents (12%) who could see the bene ts and

positive outcomes of the transition away from coal. Although the establishment of an authority to guide the transition was a possibility, the development of a renewable energy hub was considered an important opportunity. Currently, it is likely that the development of a renewable energy hub in UHV will move forward, as AGL and the Liddell Innovation Project have submitted a grant application.

Before a successful transition can take place, however, the "underlying culture of ignorance" associated with a lack of understanding about toxic chemicals and health, as well as the lack of knowledge about renewable energy, needs to be addressed through investment in community health education, renewable energy education, and educational support to provide retraining opportunities for power workers.

Ruth, L.A. (2001). Advanced coal-fired power plants. Journal of Energy Resources Technology, 123(1), 4–9. https://doi.org/10.1115/1.1348270

Slezak, M. (2016, November 7). Australian unions call for 'just transition' from coal-generated electricity. The Guardian. https://www.theguardian.com/australia-news/2016/nov/08/australian-unions-call-for-just-transition-from-coal-generated-electricity

Slezak, M. (2018, October 12). Climate target set by IPCC requires 12 Australian coal-fired power stations to close: Parliamentary Library report. ABC NEWS. https://www.abc.net.au/news/2018-10-13/coal-power-stations-needed-to-close-to-meet-ipcc-target-report/10368194

Smith, K.R., Frumkin, H., Balakrishnan, K., Butler, C.D., Chafe, Z.A., Fairlie, I., Kinney, P., Kjellstrom, T., Mauzerall, D.L., McKone, T.E., McMichael, A.J., & Schneider, M. (2013). Energy and human health. Annual Review of Public Health, 34, 159–188. https://doi.org/10.1146/annurev-publhealth-031912-114404

Streck, C., Keenlyside, P., & von Unger, M. (2016). The Paris Agreement: A new beginning. Journal for European Environmental & Planning Law, 13(1), 3–29. https://doi.org/10.1163/18760104-01301002

Sweeney, S., & Treat, J. (2018). Trade unions and just transition: The search for a transformative politics [Working Paper No. 11]. Trade Unions for Energy Democracy. http://unionsforenergydemocracy.org/wp-content/uploads/2018/04/TUED-Working-Paper-11.pdf

Thomas, N., Hook, L., & Tighe, C. (2019, September 30). How Britain ended its coal addiction. Financial Times. https://www.ft.com/content/a05d1dd4-dddd-11e9-9743-db5a370481bc

U.S. Energy Information Administration. (2020, May 11). U.S. coal-fired electricity generation in 2019 falls to 42-year low. https://www.eia.gov/todayinenergy/detail.php?id=43675

Venners, S.A., Wang, B., Xu, Z., Schlatter, W., Wang, L., & Xu, X. (2003). Particulate matter, sulfur dioxide, and daily mortality in Chongqing, China.

` DIRECT FROM AEHAP

students. In many ways, it felt like March 2020 was stuck on repeat.

Hurdles in higher education will continue to be encountered as a result of the pandemic; however, environmental health programs have the opportunity to re ect on their experiences from the past year and identify opportunities allowing for the relevance and value of environmental health to be demonstrated during the pandemic and beyond. We owe it to the students of our programs, whose unbelievable exibility and resiliency during the pandemic demonstrates their commitment to the environmental health eld. In this column, we re ect on the lessons learned from the spring and summer 2020 terms and explore opportunities to enhance the means

8ld. In

The COVID-19 pandemic has emphasized the universal importance of and need for environmental health. As a result of the pandemic, institutions of higher education have faced unprecedented changes. Decisions to keep students, faculty, and staff healthy did not come without unintended consequences as traditional means of teaching and learning were disrupted. Like faculty, students were thrust into the virtual classroom with little to no time to prepare and reported that the mid-semester transition negatively affected the overall quality of courses, which were viewed as less engaging and less instilling of learning (Garris & Fleck, 2020).

Academia collectively limped over the nish line of spring 2020 still facing a future of uncertainties. Environmental health programs scrambled to coordinate internships and meet degree requirement needs throughout the summer months in light of stay-at-home orders and nationwide emergencies. As educators, we grappled with pedagogic uncertainties surrounding the 2020–2021 academic year and anticipated the forms our classrooms could take as a result of the pandemic. The timing of instructional modality announcements throughout higher education varied, leaving some programs little time to solidify course designs and acquire the resources necessary to educate and engage

conferencing and live-streaming without all students having adequate access, we could be exasperating the inequalities we are trying to x (Gannon, 2020). While students did not begin the spring 2020 term anticipating online learning and overall student perspectives were negative, they appreciated the exibility afforded by online learning and research suggests that student acceptance and views of online learning might improve with familiarity, ideal instruction, and instructor

during the COVID-19 pandemic. Scholarship of Teaching and Learning in Psychology. Advance online publication. http://dx.doi.org/10.1037/stl0000229

Goodman, J., Melkers, J., & Pallais, A. (2019). Can online delivery increase access to education? Journal of Labor Economics, 37(1). https://doi.org/10.1086/698895

Govindarajan, V., & Srivastava, A.. (2020, March 31). What the shift to virtual learning could mean for the future of higher ed. Harvard Business Review. https://hbr.org/2020/03/what-the-shift-to-virtual-learning-could-mean-for-the-future-of-higher-ed

Smith, C.L., Coleman, S.K, & Ferrier, C. (2019). Employer and work-based student perceptions of virtual laboratory teaching and assessment resources. Work Based Learning e-Journal, 8(1), 53–70.

Supiano, B. (2020, April 23). Why is Zoom so exhausting?The Chronicle of Higher Education. https://www.chronicle.com/article/why-is-zoom-so-exhausting/

Baylor University
Waco, TX
Bryan W. Brooks, MS, PhD
bryan_brooks@baylor.edu

Benedict College
Columbia, SC
Milton A. Morris, MPH, PhD
morrism@benedict.edu

Boise State University
Boise, ID
Kim Rauscher, MA, ScD
kimberlyrauscher@boisestate.edu

California State University at Northridge †
Northridge, CA
Nola Kennedy, PhD
nola.kennedy@csun.edu

California State University at San Bernardino
San Bernardino, CA
Lal S. Mian, PhD
lmian@csusb.edu

Central Michigan University
Mount Pleasant, MI
Rebecca Uzarski, PhD
uzars2rl@cmich.edu

Colorado State University
Fort Collins, CO
Joshua Schaeffer, PhD, CIH
joshua.schaeffer@colostate.edu

East Carolina University †
Greenville, NC
William Hill (undergraduate)
hillw@ecu.edu
Stephanie Richards, PhD (graduate)
richardss@ecu.edu

East Central University
Ada, OK
Michael Bay, PhD
mbay@ecok.edu

East Tennessee State University †
Johnson City, TN
Kurt Maier, MS, PhD
maier@etsu.edu

Eastern Kentucky University †
Richmond, KY
Vonia Grabeel, MPH, RS
vonia.grabeel@eku.edu

Fort Valley State University ††
Fort Valley, GA
Oreta Samples, PhD
sampleso@fvsu.edu

Illinois State University
Normal, IL
Guang Jin, PhD, PE
gjin@ilstu.edu

Indiana University–Purdue University Indianapolis
Indianapolis, IN
Max Moreno, MEM, PhD
mmorenom@iu.edu

Mississippi Valley State University †
Itta Bena, MS
Swatantra Kethireddy, PhD
swatantra.kethireddy@mvsu.edu

Missouri Southern State University
Joplin, MO
Michael Fletcher, MS, PhD
etcher-m@mssu.edu

Montana State University
Bozeman, MT
Seth Walk, PhD
seth.walk@montana.edu
Mari Eggers, PhD
mari.eggers@montana.edu

North Carolina Central University
Durham, NC
John Bang, MD, PhD
jjbang@nccu.edu

Ohio University
Athens, OH
Michele Morrone, PhD
morrone@ohio.edu

Old Dominion University
Norfolk, VA
Sean Banaee, PhD
sbanaee@odu.edu

State University of New York, College of Environmental Science and Forestry at Syracuse
Syracuse, NY
Lee Newman, PhD
lanewman@esf.edu

Texas Southern University
Houston, TX
Zivar Youse pour, PhD
zivar.youse pour@tsu.edu

The University of Findlay †
Findlay, OH
Timothy Murphy, PhD
murphy@ ndlay.edu

Timothy477H, R72(undergradurphy Univ6.444
iD

S

` DIRECT FROM CDC ENVIRONMENTAL HEALTH SERVICES

In 2000, the Centers for Disease Control and Prevention (CDC), recognizing the important role environmental health programs play in food safety, funded a new cooperative agreement program on retail food safety called the Environmental Health Specialists Network (EHS-Net, pronounced S-Net). EHS-Net is a network of environmental health programs in state and local health departments focused on understanding how retail food service establishment policies and practices contribute to foodborne illness and outbreaks. EHS-Net staff collaborate closely with their counterparts in epidemiology and laboratory programs, and with CDC, the Food and Drug Administration (FDA), and the U.S. Department of Agriculture (USDA) Food Safety and Inspection Service (FSIS). EHS-Net staff are experienced in food safety and uniquely positioned to collect high-quality data on food safety policies and practices.

In its 20-year history, EHS-Net has conducted 15 retail food safety studies. These studies focused on restaurants because over one half of foodborne outbreaks are linked with restaurants (e.g., sit-down, fast food, deli) (Centers for Disease Control and Prevention [CDC], 2019). These studies, based primarily on data collection from observations of and interviews with restaurant staff, resulted in 50 scienti c articles and 25 plain language summaries with key ndings and recommendations (www.cdc.gov/nceh/ehs/ehsnet/publications/index.htm). A recent review of these ndings revealed key restaurant actions linked with food safety: having/adopting procedures to minimize food safety risks, training staff on those procedures, and monitoring to ensure procedures are followed (Food and Drug Administration, 2018).

Food Safety Procedures

EHS-Net found links between food safety procedures (e.g., policies, plans) and food safety.

- Restaurants with written slicer cleaning policies cleaned their food slicers more often (Brown et al., 2016).
- Workers in restaurants with a staf ng plan for when workers couldn't come to work and with an ill worker policy were less likely to have worked while ill with foodborne illness symptoms (Sumner et al., 2011).
- Restaurants with a cleaning policy had smaller norovirus outbreaks (Hoover et al., 2020).
- Restaurants with a date-marking policy practiced proper date-marking more often (Brown et al., 2021).
- Workers in restaurants with a policy prohibiting bare-hand contact with ready-to-eat food had less frequent behaviors that could lead to pathogen cross-contamination (Masters et al., 2018).

Brown, L.G., Hoover, E.R., Faw, B.V., Hedeen, N.K., Nicholas, D., Wong, M.R., Shepherd, C., Gallagher, D.L., & Kause, J.R. (2018). Food safety practices linked with proper refrigerator temperatures in retail delis. Foodborne Pathogens and Disease, 15(5), 300–307. https://doi.org/10.1089/fpd.2017.2358

Brown, gherong, , E.R.,cholas, N.K.8 (d,)Tf 0.0 Tc 0.063 Tw -1.2 Td [(N.D(, B.V)179 Ner)6969.8 25.18 (, F.K.8 (d,449 ( )]TJ -& Kau6e, Res i

DAVIS CALVIN WAGNER SANITARIAN A

Nominations for this award are open to all AAS diplomates who:

1. Exhibit resourcefulness and dedication in promoting the improvement of the public's health through the application of environmental and public health practices.
2. Demonstrate professionalism, administrative and technical skills, and competence in applying such skills to raise the level of environmental health.
3. Continue to improve through involvement in continuing education type programs to keep abreast of new developments in environmental and public health.
4. Are of such excellence to merit AAS recognition.

NOMINATIONS MUST BE RECEIVED BY APRIL 15, 2021.

Nomination packages should be e-mailed to Dr. Robert WPowitz at powitz@sanitarian.com

Files should be in Word or PDF format.

For more information about the award nomination, eligibility, and the evaluation process, as well as previous recipients of the award, please visit www.sanitarians.org/awards.

Don’t Miss This Opportunity!

Applications for the 2021 National Environmental Health Association/American Academy of Sanitarians (NEHA/AAS) Scholarship Program are now being accepted.

Undergraduate and graduate students enrolled in an accredited college or university with a dedicated curriculum in environmental health sciences are encouraged to apply.

Nomination deadline is March 31, 2021.

For eligibility information and to apply, visit www.neha.org/scholarship.

JEH QUIZ

1. In the absence of effective vaccines, vector control is the primary means of controlling the spread of arboviruses.
   a. True.
   b. False.
2. The ef cacy of pyriproxyfen autodissemination stations was assessed for Aedes albopictus and showed that Ae. albopictus carrying



2021

The Walter S. Mangold Award recognizes an individual for extraordinary achievement in environmental health. Since 1956, this award acknowledges the brightest and best in the profession. NEHA is currently accepting nominations for this award by an af liate in good standing or by any ve NEHA members, regardless of their af liation.

The Mangold is NEHA's most prestigious award and while it recognizes an individual, it also honors an entire profession for its skill, knowledge, and commitment to public health.

Nomination deadline is March 15, 2021.

RESOURCE CORNER

Resource Corner highlights different resources the National Environmental Health Association (NEHA) has available to meet your education and training needs. These resources provide you with information and knowledge to advance your professional development. Visit NEHA's online Bookstore for additional information about these and many other pertinent resources!





SPECIAL LISTING

The National Environmental Health Association (NEHA) Board of Directors includes nationally elected officers and regional vice-presidents. Affiliate presidents (or appointed representatives) comprise the Affiliate Presidents Council. Technical advisors, the executive director, and all past presidents of the association are ex-officio council members. This list is current as of press time.

Rachelle Blackham, MPH, LEHS
Region 3
Vice-President

Kim Carlton, MPH, REHS/RS, CFOI
Region 4
Vice-President

National Officers

www.neha.org/national-officers

President—Sandra Long, REHS, RS
President@neha.org

President-Elect—Roy Kroeger, REHS
roykehs@laramiecounty.com

First Vice-President—D. Gary Brown, DrPH, CIH, RS, DAAS
FirstVicePresident@neha.org

Second Vice-President—Tom Butts, MSc, REHS
SecondVicePresident@neha.org

Immediate Past-President—Priscilla Oliver, PhD
ImmediatePastPresident@neha.org

Regional Vice-Presidents

www.neha.org/RVPs

Region 1—Frank Brown, MBA, REHS/RS
Region1RVP@neha.org
Alaska, Idaho, Oregon, and Washington. Term expires 2023.

Region 2—Michele DiMaggio, REHS
Region2RVP@neha.org
Arizona, California, Hawaii, and Nevada. Term expires 2021.

Region 3—Rachelle Blackham, MPH, LEHS
Region3RVP@neha.org
Colorado, Montana, Utah, Wyoming, and members residing outside of the U.S (except members of the U.S. armed services). Term expires 2021.

Region 4—Kim Carlton, MPH, REHS/RS, CFOI
Region4RVP@neha.org
Iowa, Minnesota, Nebraska, North Dakota, South Dakota, and Wisconsin. Term expires 2022.

Region 5—Traci (Slowinski) Michelson, MS, REHS, CP-FS
Region5RVP@neha.org
Arkansas, Kansas, Louisiana, Missouri, New Mexico, Oklahoma, and Texas. Term expires 2023.

Region 6—Nichole Lemin, MS, MEP, RS/REHS
Region6RVP@neha.org
Illinois, Indiana, Kentucky, Michigan, and Ohio.
Term expires 2022.

Region 7—Tim Hatch, MPA, REHS
Region7RVP@neha.org
Alabama, Florida, Georgia, Mississippi, North Carolina,



and my dedication to constantly lead by example, innovation,



Region 2

Michele DiMaggio, REHS

My name is Michele R. DiMaggio and I was recently appointed as the National Environmental Health Association (NEHA) Regional 2 Vice-President in December 2019.

Over the past 25 years, I have gained experience as an Agricultural Biologist, Public Health Biologist, and as a Vector Control Technician. Since 2001, I have been a Registered Environmental Health Specialist working for Contra Costa County, Environmental Health Services. I am currently a Supervising Environmental Health Specialist managing the following programs: Retail Food, Temporary Food Facilities, Certified Farmer's Markets, New Employee Training and Standardization, and Schools. I also co-manage my division's Outbreak Response and Recall Team. Additionally, I participate as the Environmental Health liaison to the Health Services Department Emergency Management Team (EMT) where I serve on the EMT's Emergency Operations Response Plan Committee and act as Chair to the Training and Recruitment Committee. During the COVID-19 epidemic, I am now humbly serving as a Dhcing asg,403 0 Td ( )Tj t.015,]TJ 0 Tc 0 Tw 274 Envitment Emers EmerOver7 TOve

FDA COVID-19 Emergency Call Center as the only environmental health of cer on a 25-member of cer team directly answering 1,000+ calls of a 20,000 call volume total which was received from all 50 states and numerous international calls with questions regarding emergency use authorization for diagnostic test kits, PPE, hand sanitizer, automated ventilators, PPE decontamination methods and various aerosol sterilization or sanitation methods. He answered many basic telemedicine questions and had numerous opportunities to utilize his bi-lingual speaking skills to answer similar telehealth inquiries.

He has been a NEHA member since 2004 the same year he earned the NEHA AAS graduate student scholarship.

He has been grateful to serve on the NEHA board of directors since 2012 to present as an elected Region 8 Vice-President encompassing the mid-Atlantic States (Wash DC, WV, VA, MD, DE, PA, and all the uniformed services).

He is an alumnus with Old Dominion University Environmental Health program and active with the ODU College of Health Sciences Advisory Board since 2018. He is an active mentor to students and young professionals.

LCDR Speckhart currently chairs a NEHA Ad Hoc committee focused on Research and Innovation. He recently co-authored a NEHA policy paper on Research and Innovation that Enhance EH Science and practice published in October 2020. He is coordinating outreach with NEHA staff to develop strategic goals to foster the associations capacity to promote and appropriately support

NEHA NEWS

NEHA Announces the Dr. Bailus Walker, Jr. Diversity and Inclusion Awareness Award

Organizations around the country are striving to increase diversity awareness and the National Environmental Health Association (NEHA) is no exception. Each NEHA member has immense value and together we can achieve common goals if we foster equality and inclusion in our association. The con icts within our society that stemmed from incidents of injustice and inequity in 2020 have prompted reactions from many individuals and organizations. One step NEHA leaders are taking to denounce racism and promote diversity is the introduction of the Dr. Bailus Walker, Jr. Diversity and Inclusion Awareness Award.

Dr. Bailus Walker, Jr. had a long career as a researcher, faculty member, and administrator in public, environmental, and occupational health. Dr. Walker was a professor of environmental and occupational medicine and toxicology in the College of Medicine, Department of Community and Family Medicine at Howard University in Washington, DC. He was a consultant to public health programs seeking accreditation. He wrote proli cally in the eld of public, environmental, and occupational health. He was a long-time member and supporter of NEHA. Many of his students, leaders, employees, and mentees speak highly of his teaching, consulting, and the examples he set. He worked in several states—Alabama, California, Massachusetts, Minnesota, New York, Oklahoma, and Washington, DC—affecting many schools of public health during his career.

Dr. Walker wrote and spoke on public health, toxicology, and diversity in the eld of environmental health. He was the recipient of numerous awards for his work, including NEHA's highest honor, the 1984 Walter S. Mangold Award. He chaired a congressional CSPAN meeting on environmental racism in 1993 that pulled in several experts on the topic and shed light on the harm to communities of color. Dr. Walker headed various state and federal government working groups, committees, and task forces. He was a past president of the American Public Health Association (APHA). APHA Executive Director Dr. Georges C. Benjamin called Dr. Walker a "true public health champion." It is a tting honor for NEHA to recognize Dr. Walker by naming the new Diversity and Inclusion Awareness Award after him. It is our hope that his endeavors in diversity and environmental health will be remembered through this award.

The Dr. Bailus Walker, Jr. Diversity and Inclusion Awareness Award will be given annually to recognize an individual or group who made signi cant achievements in the development and enhancement of a more culturally diverse, inclusive, and competent environment. The quali ed individual or group will demonstrate diversity in performance in their organization and community. The application period will be open from January 15–April 15, 2021. Eligibility requirements and the award application can be found at www.neha.org/about-neha/awards/dr-bailus-walker-jr-diversity-and-inclusion-awareness-award.

NEHA Thanks the Colorado Gives Campaign Supporters

Thank you for supporting the NEHA/AAS Scholarship during NEHA's rst Colorado Gives campaign. Colorado Gives Day brought in a record number of donations made to the NEHA/AAS Scholarship fund in one single day, as well as a wealth of rst-time supporters. We are tremendously thankful for your generosity and dedication to NEHA and the NEHA/AAS Scholarship fund.

munity. NEHA is a trusted source that can speak directly about the issues that matter to you. Here are some of the ways that NEHA's Government Affairs is working to keep you informed:

Your Insider in Government Affairs Blog: Stay informed on critical and timely issues through this regularly updated blog at www.neha.org/government-affairs/your-insider-government-affairs-blog.

Government Affairs Webinar Series: If you missed any of our recent Government Affairs webinars, you can view them anytime on our website at www.neha.org/node/61387. Past webinars include: "Environmental Health Priorities at CDC, FDA, and EPA Under the Biden Administration;" Environmental Health Within the New Administration and Congress;" "Bridging the Gaps Between Environmental Quality and Public Health;" and "Food Safety Legislation Now and Future."

Position Statements and Declarations: These documents guide NEHA's efforts on public policy, directing our response to speci issues affecting environmental health. Issues covered include COVID-19 vaccination, research and innovation in environmental health.8 ssueslence, clir te change, fTJ s Legi8 (, dir)1to t wwwe

NEHA, in partnership with the Agency for Toxic

3. Human Capital: We, along with public health nurses, are the backbone of public health. The reason we were called upon to work on the opioid epidemic and COVID-19 pandemic in such large numbers is because we are highly trained professionals who are grounded in the sciences and accompanied by a high degree of relational trust throughout our communities. To be clear, on most days, when people think of public health, it is environmental health with which they identify. The postpandemic universe should emphasize a renewed alignment between us and public health nurses.
4. Physical Assets: The visible symbols of public health are generally environmental health. The Prop 65 signage in California, warnings against swimming in unhealthy surface waters, and restaurant grades are among the visible manifestations of

Kigali is a study in contrasts. The city streets are clean and virtually free of abandoned and randomly discarded trash. For those of you interested in microplastics, Rwanda began regulating plastics in 2008, a law that applies to the production and importation of most plastics, including shopping bags. The country takes this environmental practice seriously. In fact, as I transited through their border in 2010, customs of cials rif ed through my luggage aiming to unearth plastic bags. They succeeded in identifying and con scating a plastic hotel laundry sack I thoughtlessly used to separate soiled from clean clothing. I was favorably impressed by this African country's environmental policy as I have chaffed at carelessly tossed shopping bags and other plastic litter in the streets of urban environments worldwide. While this policy is a beacon of hope on a polluted planet, Kigali is also home of the Genocide Memorial, a grim reminder of humanity's most despicable and abhorrent behavior.

I visited the memorial one Saturday afternoon after spending a couple weeks ensconced in nearby Goma. While the experience of working in the Democratic Republic of the Congo was sobering, the memorial left me a changed person. The facility faithfully attempted to portray the events that led to the death of an estimated 800,000 people during the 1994 genocide. It is also a place of learning and remembrance that documents the worst of human predispositions centered on the politics of conspiracy, revenge, and retribution. I sat in silence for the better part of one hour in the memorial's garden re ecting on the horrors of East Timor, Srebrenica, and Darfur, all of which occurred in my lifetime.

I have thought about my experience in Rwanda and sought to identify what we can learn from it and its recovery from the 1994 genocide. Are there lessons we might apply to the healing necessary for us in the postpandemic U.S.? Before you rush to judgement, think for a moment. The U.S. is home to approximately 4% of the global population and yet we represented roughly 25% of the global COVID-19 case load. Our mortality experience is equally depressing. An estimated 18–20% of global COVID-19 deaths occurred in the U.S. For us to meander into the future in the absence of re ection would be unconscionable. Where do we as a nation go from here and what is our role as environmental health professionals in our nation's journey into recovery?

I offer four principles that might frame out our next steps.

1. Social Capital: I have spoken at af liate meetings on how our profession offers three unique characteristics that make us exceptionally valuable to the public health enterprise. Those are access, axis, and affect. In aggregate, we know and understand our communities and their various subcultures, commercial enterprises, and attendant risks. We should sustainably and permanently scale-up our professional relationships with the clinical and social service professions over time and share with them the valuable intelligence we possess on how our communities function. With few exceptions, health of cials who have dominated

Thank you to all who’ve tirelessly pursued public health and safety since COVID

Can your data management system optimize and map your inspector's daily schedule?

Enable your inspectors to get the most out of their day with HealthSpace. Learn more by visiting

info.gethealthspace.com/NEHA